

FIRST CASE OF FERMAT'S LAST THEOREM

Joseph Amal Nathan

Reactor Physics Design Division, Bhabha Atomic Research Centre, Mumbai-400085, India

email:josephan@magnum.barc.ernet.in

Abstract: In this paper two conjectures are proposed based on which we can prove the first case of Fermat's Last Theorem (FLT) for all primes $p \equiv -1 \pmod{6}$. With Pollaczek's result [1] and the conjectures the first case of FLT can be proved for all primes greater than 3. With a computer Conjecture1 was verified to be true for primes ≤ 2437 and Conjecture2 for primes ≤ 100003 .

Fermat's Last Theorem (FLT): *The equation $x^l + y^l = z^l$ with integral $l > 2$, has no solution in positive integers x, y, z .*

There is no loss in generality if x, y, z are pairwise prime and, if $l = 4$ and all odd primes only. Let p denote a prime greater than 3. The results presented here are when $p \nmid xyz$, known as 'first case of FLT'. We will denote greatest common divisor of integers $\alpha, \beta, \gamma, \dots$ by $(\alpha, \beta, \gamma, \dots)$.

Let us define a *Polynomial* $F_n(x, y) = (x+y)^n - x^n - y^n$ for any positive odd integer n . Expanding $(x+y)^n$ and collecting terms having same binomial coefficient we get,

$$F_n(x, y) = xy(x+y) \left[\sum_{i=1}^{(n-1)/2} {}^nC_i x^{i-1} y^{i-1} \left(\sum_{j=0}^{n-1-2i} (-1)^j x^{n-2i-j-1} y^j \right) \right]. \quad (1)$$

We will show $F_n(x, y)$ can be written in terms of x, y and $(x+y)$,

$$F_n(x, y) = xy(x+y) \sum_{i=1}^{(n-1)/2} W_{i-1} x^{i-1} y^{i-1} (x+y)^{n-2i-1}, \quad (2)$$

where, W_j , for $j = 0, 1, \dots, \frac{n-3}{2}$, are integers. Comparing (1) and (2) we get,

$$W_j = {}^nC_{j+1} - \sum_{k=1}^j \left[(-1)^{j+k} {}^nC_k + {}^{n-2j+2k-3}C_k W_{j-k} \right], \quad (3)$$

which shows all W_j 's are integers. We see $W_0 = n$ and evaluating the last coefficient,

$$\left. \frac{F_n(x, y)}{x+y} \right|_{x=-y} = (-1)^{\frac{n-1}{2}} W_{\frac{n-3}{2}} x^{n-1} = \left[(x+y)^{n-1} - \left(\frac{x^n + y^n}{x+y} \right) \right] \Big|_{x=-y} = -nx^{n-1},$$

we get

$$W_{\frac{n-3}{2}} = (-1)^{\frac{n-3}{2}} n. \quad (4)$$

Lemma1: Given nonzero integers x, y such that $(x, y) = 1$ for any positive odd integer n ,

$$\left(x + y, \frac{x^n + y^n}{x + y}\right) = (x + y, n).$$

Let $(x + y, n) = g$.

Proof 1. From (2) and (4), $\frac{F_n(x, y)}{g(x + y)} = \frac{(x + y)}{g} \left[\sum_{i=1}^{(n-3)/2} W_{i-1} x^i y^i (x + y)^{n-2i-2} \right] - \frac{n}{g} (-xy)^{\frac{n-1}{2}}$, any divisor of $\frac{(x + y)}{g}$ will divide all terms except last, giving $\left(\frac{x + y}{g}, \frac{F_n(x, y)}{g(x + y)}\right) = 1$. So $\left(\frac{x + y}{g}, \frac{x^n + y^n}{g(x + y)}\right) = 1$.

Proof 2. Since, $\frac{x^n + y^n}{g(x + y)} = \frac{(x + y)}{g} \left[\sum_{i=1}^{n-1} (-1)^{i-1} i x^{n-i-1} y^{i-1} \right] + \frac{n}{g} y^{n-1}$, any divisor of $\frac{(x + y)}{g}$ will divide all terms except last. Hence $\left(\frac{x + y}{g}, \frac{x^n + y^n}{g(x + y)}\right) = 1$. $\square$

Now let $n = p$, then from (3), $w_j = W_j/p$, for $j = 0, 1, \dots, \frac{n-3}{2}$, are integers. Define *Polynomials* $G_p(x, y) = \frac{F_p(x, y)}{p x y (x + y)}$, and $f_{x, y} = x^2 + xy + y^2$. We will now show, $f_{x, y}^\epsilon$ is a factor of $G_p(x, y)$ for any prime $p > 3$, when exponent $\epsilon = 1$. We will also show $\epsilon = 2$ if and only if $p \equiv 1 \pmod{6}$ and $\epsilon \leq 2$. Solving for x in $f_{x, y} = 0$, gives $x = -ye^{\pm \frac{i\pi}{3}}$. Now to show $f_{x, y}^\epsilon \mid G_p(x, y)$ it is sufficient to show $\frac{F_p(x, y)}{f_{x, y}^{\epsilon-1}} \Big|_{x=-ye^{\pm \frac{i\pi}{3}}} = 0$. When $\epsilon = 1$ substituting for x in $F_p(x, y) = (x + y)^n - x^n - y^n$, we get

$$F_p(x, y) \Big|_{x=-ye^{\pm \frac{i\pi}{3}}} = y^p \left[2 \cos \frac{p\pi}{3} - 1 \right],$$

which is zero for any prime $p \equiv \pm 1 \pmod{6}$. Since $F_p(x, y)/f_{x, y}^{\epsilon-1}$ at $x = -ye^{\pm \frac{i\pi}{3}}$ is indeterminant for $\epsilon > 1$, applying L'Hospital's rule when $\epsilon = 2$ we get,

$$\frac{F_p(x, y)}{f_{x, y}} \Big|_{x=-ye^{\pm \frac{i\pi}{3}}} = \frac{[F_p(x, y)]'}{[f_{x, y}]'} \Big|_{x=-ye^{\pm \frac{i\pi}{3}}} = \mp 2ip y^{p-2} \left[\frac{\sin \frac{(p-1)\pi}{3}}{1 - 2 e^{\pm \frac{i\pi}{3}}} \right],$$

which is zero if and only if $p \equiv 1 \pmod{6}$. When $\epsilon = 3$ we get,

$$\frac{F_p(x, y)}{f_{x, y}^2} \Big|_{x=-ye^{\pm \frac{i\pi}{3}}} = \frac{[F_p(x, y)]''}{[f_{x, y}]''} \Big|_{x=-ye^{\pm \frac{i\pi}{3}}} = p(p-1)y^{p-4} \left[\frac{\cos \frac{(p-2)\pi}{3}}{\left(1 - 2 e^{\pm \frac{i\pi}{3}}\right)^2} \right],$$

which is not zero for $p \equiv \pm 1 \pmod{6}$. So $\epsilon \leq 2$ and $\epsilon = 1$ or 2 for primes $p \equiv -1$ or $+1 \pmod{6}$ respectively.

Lemma2: If the first case of FLT is false, then $p^2 \mid G_p(x, y)$ for primes $p \geq 3$.

Since first case of FLT is false we have nonzero pairwise prime integers x, y, z satisfying equation $x^p + y^p = z^p$. Define $m = x + y - z$, from Fermat's little theorem $p \mid m$. Let $p^\lambda \mid m$ first we will show $\lambda \geq 2$. Since $p \nmid xyz$ we have $(x + y, p) = 1$ and from Lemma1

$$(x + y) = a^p; \quad (z - x) = b^p; \quad (z - y) = c^p, \quad (5)$$

$$\frac{x^p + y^p}{x + y} = r^p; \quad \frac{z^p - x^p}{z - x} = s^p; \quad \frac{z^p - y^p}{z - y} = t^p, \quad (6)$$

$$z = ar; \quad y = bs; \quad x = ct, \quad (7)$$

where a, b, c, r, s, t are positive integers > 1 and $p \nmid a, b, c, r, s, t$. Substituting (5) and (7) in $m = x + y - z$ we get, $m = a(a^{p-1} - r) = b(s - b^{p-1}) = c(t - c^{p-1})$. Since $p \mid m$, we have $(a^{p-1})^p \equiv r^p \pmod{p^2} \Rightarrow r^p \equiv 1 \pmod{p^2}$ and substituting for r^p from (6) we get,

$$z^p \equiv (x + y) \pmod{p^2}. \quad (8)$$

Similarly

$$y^p \equiv (z - x) \pmod{p^2}, \quad (9)$$

$$x^p \equiv (z - y) \pmod{p^2}. \quad (10)$$

Taking (8)-(9)-(10), we get

$$m \equiv 0 \pmod{p^2}, \quad (11)$$

so $\lambda \geq 2$. From (8)+(9)+(10), (8)+(9)-(10) and (8)-(9)+(10) we get $2z^p \equiv 2z \pmod{p^2}$, $2y^p \equiv 2y \pmod{p^2}$ and $2x^p \equiv 2x \pmod{p^2}$ respectively, which gives

$$z^{p-1} \equiv 1 \pmod{p^2}; \quad y^{p-1} \equiv 1 \pmod{p^2}; \quad x^{p-1} \equiv 1 \pmod{p^2}. \quad (12)$$

Now from (11) we have $(x + y)^p - z^p \equiv 0 \pmod{p^3} \Rightarrow p^2 \mid xy(x + y)G_p(x, y)$. Since $p \nmid xyz \Rightarrow p^2 \mid G_p(x, y)$ $\square$

Corollary1: For prime 3 the first case of FLT is true

Proof. Assume the first case of FLT is false for $p = 3$. We have $G_3(x, y) = \frac{(x+y)^3 - x^3 - y^3}{3xy(x+y)} = 1$. But from Lemma2, $3^2 \mid G_3(x, y)$, which is not possible. $\square$

Lemma3: For any nonzero positive coprime integers x, y there is no prime $\xi \equiv -1 \pmod{6}$ such that $\xi \mid (x^2 + xy + y^2)$.

Proof. We have $\xi + 1 = 6\tau$, where τ is a positive integer ≥ 1 . Let $x + y = \eta$. Assume $f_{x,y} = \eta^2 - xy \equiv 0 \pmod{\xi}$. Since $(x, y) = 1$ we get $(x, \xi) = (y, \xi) = (\eta, \xi) = (x, \eta) = (y, \eta) = 1$ and the following congruences,

$$x^2 \equiv -y\eta \pmod{\xi}; \quad y^2 \equiv -\eta x \pmod{\xi}; \quad \eta^2 \equiv xy \pmod{\xi}, \quad (13)$$

$$x^2 + y^2 + \eta^2 \equiv 0 \pmod{\xi}. \quad (14)$$

From (13) we get

$$x^3 \equiv y^3 \equiv -\eta^3 \pmod{\xi}. \quad (15)$$

Using Fermat's little theorem and (15) in (14) we have,

$$x^2 + y^2 + \eta^2 \equiv x^{\xi+1} + y^{\xi+1} + \eta^{\xi+1} \equiv (x^{3\tau})^2 + (y^{3\tau})^2 + (\eta^{3\tau})^2 \equiv 3x^{6\tau} \equiv 0 \pmod{\xi},$$

which is contradictory to $\xi \nmid xy\eta$. □

Corollary2: *For prime 5 the first case of FLT is true*

Proof. Assume the first case of FLT is false for $p = 5$. We have $G_5(x, y) = x^2 + xy + y^2$ and $5 \equiv -1 \pmod{6}$. But from Lemma2 we have $5^2 \mid G_5(x, y)$, contradicting Lemma3. □

Define $H_p(x, y) = G_p(x, y)/f_{x,y}^\epsilon = \frac{F_p(x,y)}{pxy(x+y)f_{x,y}^\epsilon}$. It is obvious that $H_p(x, y)$ can be defined only for primes $p > 3$. Now we propose the following *conjecture*.

Conjecture1: *For nonzero coprime positive integers x, y and primes $p > 3$, $H_p(x, y)$ is not divisible by p^2 .* Here we note that there is no condition on x and y except $(x, y) = 1$.

Proposition1: *The first case of FLT is true for primes $p \equiv -1 \pmod{6}$.* We will now show how Conjecture1 can be used to prove Proposition1. Assume the first case of FLT is false, then from Lemma2 $p^2 \mid G_p(x, y)$. But from Lemma3 we see that $p \nmid f_{x,y}$. So $p^2 \mid H_p(x, y)$ contradicting Conjecture1.

Pollaczek's result [1]: *If the non-zero integers x, y, z satisfy equation $x^p + y^p = z^p$, $(x, y, z) = 1$, for prime $p > 3$, then $x^2 + xy + y^2$ is non-divisible by p .*

Proposition2: *The first case of FLT is true for any prime $p > 3$.* Again we will show how Conjecture1 can be used to prove Proposition2. As we have already seen if the first case of FLT is false then $p^2 \mid G_p(x, y)$. But from Pollaczek's result we see that $p \nmid f_{x,y}$. So $p^2 \mid H_p(x, y)$ contradicting Conjecture1.

We propose a weaker *conjecture* which can be easily verified without altering the propositions.

Conjecture2: *For any prime $p > 3$ assume that there exist pairwise relatively prime nonzero integers x, y, z , $x^p + y^p = z^p$. Then $H_p(x, y)$ is not divisible by p^2 .*

Since $H_5(x, y) = H_7(x, y) = 1$, the conjectures are true for primes 5 and 7. For primes $p > 7$ using a simple computer programme in Mathematica 4.0 the *conjectures* were being verified. Since $F_p(x, y)$ and $xy(x+y)f_{x,y}^\epsilon$ are symmetric polynomials, $H_p(x, y) = \frac{F_p(x,y)}{pxy(x+y)f_{x,y}^\epsilon}$ is also a symmetric polynomial. So $(y^{-1})^{p-2\epsilon-3} H_p(x, y) \equiv H_p(X, 1) \pmod{p \text{ or } p^2}$, where $X \equiv y^{-1}x \pmod{p \text{ or } p^2}$ and $y^{-1}y \equiv 1 \pmod{p \text{ or } p^2}$. Due to the property $H_p(X, 1) = H_p(-X - 1, 1)$ the number of residues, of p or p^2 , for the verification of the conjectures are reduced to half. For most primes p , we found, $p \nmid H_p(X, 1)$ and the number of residues required for both conjectures is $(p-1)/2$. But when $p \mid H_p(X, 1)$ verification of Conjecture1 takes longer than Conjecture2, since the number of residues of p^2 for Conjecture1 is $p(p-1)/2$ and for Conjecture2, because of condition from (12), $X^{p-1} \equiv 1 \pmod{p^2}$, it is only $(p-1)/2$. With the computer Conjecture1 was verified to be true for primes ≤ 2437 and Conjecture2 for primes ≤ 100003 . From Corollary1, the first case of FLT is true for prime 3. So with Conjecture2 and Pollaczek's result, **the first case of FLT is true for all primes ≤ 100003** . We conclude this paper with a *corollary*.

Corollary3: *For any prime p , $2^{p-1} - 1$ is not divisible by p^3 .* This corollary can be easily checked for primes 2 and 3. For primes $p > 3$, we have from Conjecture1 or Conjecture2 $p^2 \nmid H_p(x, y)$ even for $x = y = 1$. But $H_p(x, y) = \frac{F_p(x,y)}{pxy(x+y)f_{x,y}^\epsilon}$ and substituting $x = y = 1$ we get,

$$p^2 \nmid \frac{2^{p-1}-1}{3^e p} \Rightarrow p^3 \nmid 2^{p-1} - 1. \quad \square$$

Acknowledgement. I am very thankful to M.A. Prasad for the discussions which helped in improvement of this paper. His suggestions in the computer programme reduced the run time considerably. I am highly indebted to him for his valuable guidance and encouragment. I thank V. Nandagopal, School of Mathematics, Tata Institute of Fundamental Research, Mumbai, for helping in executing the programme.

REFERENCES

1. Pollaczek, F., *Über den großen Fermatschen Satz*, Sitzungsber. Akad. Wiss. Wein. Math.-Natur. Kl. IIa, **126** (1917), 45-59.